



Influence of Macrocolumnar EEG on Ca Waves

Corresponding Author:

Prof. Lester Ingber,
R&D, Lester Ingber Research, 97520 - United States of America

Submitting Author:

Prof. Lester Ingber,
R&D, Lester Ingber Research, 97520 - United States of America

Article ID: WMC003701

Article Type: Research articles

Submitted on: 14-Sep-2012, 01:33:48 PM GMT **Published on:** 15-Sep-2012, 05:26:38 PM GMT

Article URL: http://www.webmedcentral.com/article_view/3701

Subject Categories: NEUROSCIENCES

Keywords: short-term memory, astrocytes, neocortical dynamics, vector potential

How to cite the article: Ingber L. Influence of Macrocolumnar EEG on Ca Waves . WebmedCentral NEUROSCIENCES 2012;3(9):WMC003701

Copyright: This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC-BY\)](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Source(s) of Funding:

None

Competing Interests:

None

Influence of Macrocolumnar EEG on Ca Waves

Author(s): Ingber L

Abstract

A “smoking gun” for explicit top-down neocortical mechanisms that directly drive bottom-up processes that describe memory, attention, etc. The top-down mechanism considered are macrocolumnar **EEG** firings in neocortex, as described by a statistical mechanics of neocortical interactions (**SMNI**), developed as a magnetic vector potential A . The bottom-up process considered are Ca^{2+} waves prominent in synaptic and extracellular processes that are considered to greatly influence neuronal firings. Here, the complimentary effects are considered, i.e., the influence of A on Ca^{2+} momentum, p . The canonical momentum of a charged particle in an electromagnetic field, $\vec{p} = \vec{p} + q\vec{A}$ (SI units), is calculated, where the charge of Ca^{2+} is $q = 2e$, e is the magnitude of the charge of an electron, valid in both classical and quantum mechanics. It is shown that A is large enough to influence p . This suggests that, instead of the common assumption that Ca^{2+} waves contribute to neuronal activity, they may in fact at times be caused by the influence of A of larger-scale **EEG**.

Smoking Gun For Top-Down Processes

There is a growing awareness of the importance of multiple scales in many physical and biological systems, including neuroscience [1], [2]. As yet, there does not seem to be any “smoking gun” for explicit top-down mechanisms that directly drive bottom-up processes that describe memory, attention, etc. Of course, there are many top-down type studies demonstrating that neuromodulator [3] and neuronal firing states, e.g., as defined by **EEG** frequencies, can modify the milieu or context of individual synaptic and neuronal activity, which is still consistent with ultimate bottom-up paradigms. However, there is a logical difference between top-down milieu as conditioned by some prior external or internal conditions, and some direct top-down processes that directly cause bottom-up interactions specific to **STM**. Here, the operative word is “cause”.

A. Magnetism Influences in Living Systems

There is a body of evidence that suggests a specific top-down mechanism for neocortical **STM** processing. An example of a direct physical mechanism that affects neuronal processing not part of “standard” sensory influences is the strong possibility of magnetic influences in birds at quantum levels of interaction [4]–[6]. It should be noted that this is just a proposed mechanism [7]. The strengths of magnetic fields in neocortex may be at a threshold to directly influence synaptic interactions with astrocytes, as proposed for long-term memory (**LTM**) [8] and short-term memory (**STM**) [9], [10]. Magnetic strengths associated by collective **EEG** activity at a columnar level gives rise to even stronger magnetic fields. Columnar excitatory and inhibitory processes largely take place in different neocortical laminae, providing possibilities for more specific mechanisms.

SMNI CONTEX

Since 1981 about 30 papers on a statistical mechanics of neocortical interactions (**SMNI**) has been detailed properties of short-term memory, long-term memory, **EEG** analyses, and other properties of neocortex [11]–[16].

This discussion compares the momentum of a Ca^{2+} ion with macrocolumnar **EEG** fields. Columnar **EEG** firings calculated by **SMNI** lead to electromagnetic fields which can be described by a vector potential 4-vector [17]. In the standard gauge, the 3-vector components of this vector potential describe magnetic fields, denoted here as A , are of interest. In this context this is referred to as the **SMNI** vector potential (**SMNI-VP**). An early discussion of **SMNI-VP** contained in a review of short-term memory as calculated by **SMNI** was not as detailed [16]. Note that gauge of A is not specified here, and this can lead to important effects especially at quantum scales [18]. Current research is directed to more detailed interactions of **SMNI-VP** firing states with Ca^{2+} waves.

This paper concerns a dipole model for collective minicolumnar oscillatory currents, corresponding to top-down signaling, flowing in ensembles of axons, not for individual neurons. The top-down signal is claimed to cause relevant effects on the surrounding milieu, but is not appropriate outside these surfaces due to strong attenuation of electrical activity. However, the vector potentials produced by these dipoles due to

axonal discharges do survive far from the axons, and this can lead to important effects at the molecular scale, e.g., in the environment of ions [19], [20].

The **SMNI** columnar probability distributions, derived from statistical aggregation of synaptic and neuronal interactions among minicolumns and macrocolumns, have established credibility at columnar scales by detailed calculations of properties of **STM**. Under conditions enhancing multiple attractors, detailed in SMNI papers with a "centering mechanism" effected by changes in background synaptic activity, multiple columnar collective firing states are developed. It must be stressed that these minicolumns are the entities which the above dipole moment is modeling. The Lagrangian of the **SMNI** distributions, although possessing multivariate nonlinear means and covariance, have functional forms similar to arguments of firing distributions of individual neurons, so that the description of the columnar dipole above is a model faithful to the standard derivation of a vector potential from an oscillating electric dipole.

Note that this is not necessarily the only or most popular description of electromagnetic influences in neocortex, which often describes dendritic presynaptic activity as inducing large scale EEG [21], or axonal firings directly affecting astrocyte processes [22]. This work is only and specifically concerned with electromagnetic fields in collective axonal firings, directly associated with columnar **STM** phenomena in **SMNI** calculations, which create vector potentials influencing ion momenta just outside minicolumnar structures.

Ca²⁺

The roles of Ca²⁺, while not completely understood, are very well appreciated as being quite important. It is likely that Ca²⁺ waves are instrumental in tripartite synaptic interactions of astrocytes and neuronal synapses [23]–[25].

A. Ca²⁺ Momentum

The momentum at issue is calculated to set the stage for comparison to the vector potential.

In neocortex, a Ca²⁺ ion with mass ^mCa = 6.6*10⁻²⁶ kg, has speed on the order of 50 μm/s [26] to 100 μm/s [25]. This gives a momentum on the order of 10⁻³⁰ kg-m/s. An estimate of molar concentrations [25], gives an estimate of a Ca²⁺ wave as comprised of tens of thousands of such ions.

VECTOR POTENTIAL OF EEG DIPOLES

The effective momentum, \vec{p} , affecting the momentum \vec{p} of a moving particle in an electromagnetic field, is understood from the canonical momentum [19], [27], [28], in SI units,

$$\vec{p} = \vec{p} + q\vec{A} \quad (1)$$

where $q = 2e$ for Ca²⁺, e is the magnitude of the charge of an electron 1.6 10⁻¹⁹ C (Coulomb), and \vec{A} is the electromagnetic vector potential. (Note that in Gaussian units $\vec{p} = \vec{p} + q\vec{A}/c$, where c is the speed of light.

\vec{p} can be used in quantum as well as in classical calculations. Eventually, quantum mechanical calculations including these effects will be performed, as it is clear that in time scales much shorter than neuronal firings Ca²⁺ wave packets spread over distances the size of typical synapses [29]. Note that gauge of \vec{A} is not specified here, and this can lead to important effects especially at quantum scales [18].

For a wire/neuron carrying a current I , measured in A = Amperes = C/s,

$$A = \mu_0/4\pi \int (dr/r I) \quad (2)$$

where the current is along a length z (a neuron), observed from a perpendicular distance s . Neglecting far-field retardation effects, this yields

$$A = \mu_0/4\pi I \log(z + (z^2 + s^2)^{1/2}/s) \quad (3)$$

Other formulae for other geometries are in texts [17]. The point here is the insensitive log dependence on distance. The estimates below assume this log factor to be of order 1. The magnetic field B derived from A ,

$$B = \vec{\nabla} \times \vec{A} \quad (4)$$

is still attenuated in the glial areas where Ca²⁺ waves exist, and its magnitude decreases as inverse distance, but A derived near the minicolumns will be used there and at further distance since it is not so attenuated. The electrical dipole for collective minicolumnar EEG derived from A is

$$E = ic \vec{\nabla} \times \vec{B} = ic \vec{\nabla} \times (\vec{\nabla} \times \vec{A}) \quad (5)$$

μ_0 , the magnetic permeability in vacuum = 4 π 10⁻⁷H/m (Henry/meter), where Henry has units of kg-m-C⁻², is

the conversion factor from electrical to mechanical variables. Near neurons,

$$\mu = 10\mu\text{0}[30], \text{ giving } \mu = 10^{-6}$$

qA can be calculated at several scales:

In studies of small ensembles of neurons [31], an electric dipole moment \mathbf{Q} is defined as $z\hat{r}$, where \hat{r} is the direction unit-vector, leading to estimates of $|\mathbf{Q}|$ for a pyramidal neuron on the order of $1 \text{ pA}\cdot\text{m} = 10^{-12} \text{ A}\cdot\text{m}$. Multiplying by 10^4 synchronous firings in a macrocolumn gives an effective dipole moment $|\mathbf{Q}| = 10^{-8} \text{ A}\cdot\text{m}$. Taking z to be $10^2 \mu\text{m} = 10^{-4} \text{ m}$ (a couple of neocortical layers) to get I , this gives an estimate $|\mathbf{qA}| = 2 \times 10^{-19} \times 10^{-6} \times 10^{-8} / 10^{-4} = 10^{-27} \text{ kg}\cdot\text{m/s}$,

Estimates at larger scales [32] give a dipole density $P = 0.1 \mu\text{A}/\text{mm}^2$. Multiplying this density by a volume of $\text{mm}^2 \times 10^2 \mu\text{m}$ (using the same estimate above for z), gives a $|\mathbf{Q}| = 10^{-9} \text{ A}\cdot\text{m}$. This is smaller than that above, due to this estimate including cancellations giving rise to scalp EEG, while the estimate above is within a macrocolumn (the focus of this study), leading to $|\mathbf{qA}| = 10^{-28} \text{ kg}\cdot\text{m/s}$.

SMNI CALCULATIONS

A. Ca^{2+} Momenta

The time dependence of Ca^{2+} wave momenta is typically calculated with simulations using code such as

NEURON [33], within multivariate differential equations describing interactions among quite a few neuronal elements and parameters. In this study, the resulting flow of Ca^{2+} wave momenta will be further determined by its interactions in \mathcal{V} , the canonical momenta which includes **A**.

B. SMNI-VP

The outline of coupling the **SMNI-VP** with Ca^{2+} waves follows. Similar to the scaling of mesoscopic columnar firings to an electric potential ϕ describe regional **EEG** that was fitted to large data sets [15], here columnar firings are scaled to describe the effective current \mathbf{I} giving rise to the vector potential \mathbf{A} ,

$$\mathbf{A} = aM^E \mathbf{r} + bM^I \mathbf{r} \text{ -----(6)}$$

where a and b are scaled to something on the order of 10^4 pA , as discussed above. M^E is the excitatory columnar firing of pyramidal neurons, and M^I is the inhibitory columnar firing of pyramidal neurons.

The influence of time-dependent Ca^{2+} waves is introduced in the post-synaptic and pre-synaptic SMNI parameters, which here also are time-dependent as

functions of changing Ca^{2+} ions.

Such parameters are present at neuronal scales and are included in microscopic **NEURON** ordinary differential equation calculations. However, as in the original development of **SMNI**, these parameters are developed to mesocolumnar scales.

For example, SMNI mesoscopic firings are described by coupled stochastic differential equations, nonlinear in the drifts and covariance in terms of M^E and M^I variables, and mesoscopic synaptic and neuronal parameters. It has been most productive to cast these coupled equations into mathematically equivalent conditional probability distributions, which are better suited to handle algebraic intricacies of their rather general nonlinear time-dependent structure, and which afford

the use of powerful derivations based on the associated variational principle, e.g., Canonical Momenta Indicators and Euler-Lagrange equations. This is all rigorously discussed and calculated in many preceding **SMNI** papers. This also required developing powerful numerical algorithms to fit these algebraic models to data [34], [35] and to develop numerical details of the propagating probability distributions using **PATHINT** [36] and **PATHTREE** [37].

C. Coupled SMNI-VP Ca^{2+} System

For several decades the stated Holy Grail of chemical, biological and biophysical research into neocortical information processing has been to reduce such neocortical phenomena into specific bottom-up molecular and smaller-scale processes [39]. Over the past three decades, with regard to short-term memory (**STM**) and long-term memory (**LTM**) phenomena, which themselves are likely components of other phenomena like attention and consciousness, the **SMNI** approach has

yielded specific details of **STM** capacity, duration and stability not present in molecular approaches, but it is clear that most molecular approaches consider it inevitable that their approaches at molecular and possibly even quantum scales will yet prove to be causal explanations of such phenomena.

The **SMNI** approach is a bottom-up aggregation from synaptic scales to columnar and regional scales of neocortex, and has been merged with larger non-invasive **EEG** scales with other colleagues – all at scales much coarser than molecular scales. As with many Crusades for some truths, other truths can be trampled. It is proposed that an SMNI vector potential (**SMNIVP**) constructed from magnetic fields induced by neuronal electrical firings, at thresholds of collective minicolumnar activity with laminar specification, can

give rise to causal top-down mechanisms that effect molecular excitatory and inhibitory processes in **STM** and **LTM**. Such a smoking gun for top-down effects awaits forensic in vivo experimental verification, requiring appreciating the necessity and due diligence of including true multiple-scale interactions across orders of magnitude in the complex neocortical environment.

While many studies have examined the influences of changes in Ca^{2+} distributions on large-scale **EEG** [40], there have not been studies examining the complimentary effects on Ca^{2+} ions at a given neuron site from **EEG**-induced magnetic fields arising from other neuron sites

Thus, a single Ca^{2+} ion can have a momentum appreciably altered in the presence of macrocolumnar **EEG** firings, and this effect is magnified when many ions in a wave are similarly affected. Therefore, large-scale top-down neocortical processing giving rise to measurable scalp **EEG** can directly influence atomic-scale bottom-up processes.

This suggests that, instead of the common assumption that Ca^{2+} waves contribute to neuronal activity, they may in fact at times be caused by the influence of A of larger-scale EEG.

Such a "smoking gun" for top-down effects awaits forensic in vivo experimental verification, requiring appreciating the necessity and due diligence of including true multiple-scale interactions across orders of magnitude in the complex neocortical environment.

Acknowledgments

I thank Paul Nunez and William Ross for verification of some experimental data.

References

1. C. Anastassiou, R. Perin, H. Markram, and C. Koch, "Ephaptic coupling of cortical neurons," *Nature Neuroscience*, vol. 14, pp. 217–223, 2011.
2. P. Nunez, R. Srinivasan, and L. Ingber, "Theoretical and experimental electrophysiology in human neocortex: Multiscale correlates of conscious experience," in *Multiscale Analysis and Nonlinear Dynamics*, M. Pesenson, Ed. New York: Wiley, 2012, p. (to be published).
3. R. Silberstein, "Neuromodulation of neocortical dynamics," in *Neocortical Dynamics and Human EEG Rhythms*, P. Nunez, Ed. New York, NY: Oxford

University Press, 1995, pp. 628–681.

4. I. Kominis, "Zeno is pro Darwin: quantum Zeno effect suppresses the dependence of radical-ion-pair reaction yields on exchange and dipolar interactions," University of Crete, Greece, Tech. Rep. arXiv:0908.0763v2 [quant-ph], 2009.
5. C. Rodgers and P. Hore, "Chemical magnetoreception in birds: The radical pair mechanism," *PNAS*, vol. 106, no. 2, pp. 353–360, 2009.
6. I. Solov'yov and K. Schulten, "Magnetoreception through cryptochrome may involve superoxide," *Biophys. J.*, vol. 96, no. 12, pp. 4804–4813, 2009.
7. S. Johnsen and K. Lohmann, "Magnetoreception in animals," *Phys. Today*, vol. 61, pp. 29–35, 2008.
8. G. Gordon, K. Iremonger, S. Kantevari, G. Ellis Davies, B. MacVicar, and J. Bains, "Astrocyte-mediated distributed plasticity at hypothalamic glutamate synapses," *Neuron*, vol. 64, pp. 391–403, 2009.
9. M. Banachlocha, "Neuromagnetic dialogue between neuronal minicolumns and astroglial network: A new approach for memory and cerebral computation," *Brain Res. Bull.*, vol. 73, pp. 21–27, 2007.
10. J. A. Pereira and F. Furlan, "Astrocytes and human cognition: Modeling information integration and modulation of neuronal activity," *Progress in Neurobiology*, vol. 92, pp. 405–420, 2010.
11. L. Ingber, "Statistical mechanics of neocortical interactions. I. Basic formulation," *Physica D*, vol. 5, pp. 83–107, 1982, <http://www.ingber.com/smni82n basic.pdf>.
12. L. Ingber, "Statistical mechanics of neocortical interactions. Dynamics of synaptic modification," *Phys. Rev. A*, vol. 28, pp. 395–416, 1983, <http://www.ingber.com/smni83n dynamics.pdf>.
13. L. Ingber, "Statistical mechanics of neocortical interactions. Derivation of short-term-memory capacity," *Phys. Rev. A*, vol. 29, pp. 3346–3358, 1984, <http://www.ingber.com/smni84n stm.pdf>.
14. L. Ingber, "Statistical mechanics of neocortical interactions: Pathintegral evolution of short-term memory," *Phys. Rev. E*, vol. 49, no. 5B, pp. 4652–4664, 1994, <http://www.ingber.com/smni94n stm.pdf>.
15. L. Ingber, "Statistical mechanics of neocortical interactions: Applications of canonical momenta indicators to electroencephalography," *Phys. Rev. E*, vol. 55, no. 4, pp. 4578–4593, 1997, <http://www.ingber.com/smni97n cmi.pdf>.
16. L. Ingber, "Columnar EEG magnetic influences on molecular development of short-term memory," in *Short-Term Memory: New Research*, G. Kalivas and S.

- Petralia, Eds. Hauppauge, NY: Nova, 2012a, pp. 37–72, Invited Paper. http://www.ingber.com/smni11n_stmn_scales.pdf.
17. J. Jackson, *Classical Electrodynamics*. New York: Wiley & Sons, 1962.
18. J. Tollaksen, Y. Aharonov, A. Casher, T. Kaufherr, and S. Nussinov, "Quantum interference experiments, modular variables and weak measurements," *New J. Phys.*, vol. 12, no. 013023, pp. 1–29, 2010.
19. R. Feynman, R. Leighton, and M. Sands, *The Feynman Lectures on Physics*. Reading, MA: Addison-Wesley, 1964.
20. G. Giuliani, "Vector potential, electromagnetic induction and 'physical meaning'," *Eur. J. Phys.*, vol. 31, no. 4, pp. 871–880, 2010.
21. P. Nunez, *Electric Fields of the Brain: The Neurophysics of EEG*. London: Oxford University Press, 1981.
22. J. McFadden, "Conscious electromagnetic field theory," *NeuroQuantology*, vol. 5, no. 3, pp. 262–270, 2007.
23. C. Agulhon, J. Petravicz, A. McMullen, E. Sweger, S. Minton, S. Taves, K. Casper, T. Fiacco, and K. McCarthy, "What is the role of astrocyte calcium in neurophysiology?" *Neuron*, vol. 59, pp. 932–946, 2008.
24. A. Araque and M. Navarrete, "Glial cells in neuronal network function," *Phil. Tran. R. Soc. B*, pp. 2375–2381, 2010.
25. W. Ross, "Understanding calcium waves and sparks in central neurons," *Nature*, vol. 13, pp. 157–168, 2012.
26. S. Bellingier, "Modeling calcium wave oscillations in astrocytes," *Neurocomputing*, vol. 65, no. 66, pp. 843–850, 2005.
27. R. Feynman, *Quantum Electrodynamics*. New York: W.A. Benjamin, 1961.
28. H. Goldstein, *Classical Mechanics*, 2nd ed. Reading, MA: Addison Wesley, 1980.
29. H. Stapp, *Mind, Matter and Quantum Mechanics*. New York: Springer-Verlag, 1993.
30. D. Georgiev, "Electric and magnetic fields inside neurons and their impact upon the cytoskeletal microtubules," *Cogprints*, U. Southampton, UK, Tech. Rep. *Cogprints Report*, 2003, <http://cogprints.org/3190/>.
31. S. Murakami and Y. Okada, "Contributions of principal neocortical neurons to magnetoencephalography and electroencephalography signals," *J. Physiol.*, vol. 575, no. 3, pp. 925–936, 2006.
32. P. Nunez and R. Srinivasan, *Electric Fields of the Brain: The Neurophysics of EEG*, 2nd Ed. London: Oxford University Press, 2006.
33. N. Carnevale and M. Hines, *The NEURON Book*. Cambridge, UK: Cambridge U Press, 2006.
34. L. Ingber, "Adaptive Simulated Annealing (ASA)," Caltech Alumni Association, Pasadena, CA, Tech. Rep. *Global optimization C-code*, 1993, <http://www.ingber.com/n#ASA-CODE>.
35. L. Ingber, "Adaptive Simulated Annealing," in *Stochastic global optimization and its applications with fuzzy adaptive simulated annealing*, J. H.A. Oliveira, A. Petraglia, L. Ingber, M. Machado, and M. Petraglia, Eds. New York: Springer, 2012b, pp. 33–61, Invited Paper. http://www.ingber.com/asa11n_options.pdf.
36. L. Ingber and P. Nunez, "Statistical mechanics of neocortical interactions: High resolution path-integral calculation of short-term memory," *Phys. Rev. E*, vol. 51, no. 5, pp. 5074–5083, 1995, http://www.ingber.com/smni95n_stm.pdf.
37. L. Ingber, C. Chen, R. Mondescu, D. Muzzall, and M. Renedo, "Probability tree algorithm for general diffusion processes," *Phys. Rev. E*, vol. 64, no. 5, pp. 056 702–056 707, 2001, http://www.ingber.com/path01n_pathtree.pdf.
38. H. Dekker, "On the most probable transition path of a general diffusion process," *Phys. Lett. A*, vol. 80, pp. 99–101, 1980.
39. M. Rabinovich, P. Varona, A. Selverston, and H. Arbaranel, "Dynamical principles in neuroscience," *Rev. Mod. Phys.*, vol. 78, no. 4, pp. 1213–1265, 2006.
40. P. Kudela, G. Bergey, and P. Franaszczuk, "Calcium involvement in regulation of neuronal bursting in disinhibited neuronal networks: Insights from calcium studies in a spherical cell model," *Biophys. J.*, vol. 97, no. 12, pp. 3065–3074, 2009.

Disclaimer

This article has been downloaded from WebmedCentral. With our unique author driven post publication peer review, contents posted on this web portal do not undergo any prepublication peer or editorial review. It is completely the responsibility of the authors to ensure not only scientific and ethical standards of the manuscript but also its grammatical accuracy. Authors must ensure that they obtain all the necessary permissions before submitting any information that requires obtaining a consent or approval from a third party. Authors should also ensure not to submit any information which they do not have the copyright of or of which they have transferred the copyrights to a third party.

Contents on WebmedCentral are purely for biomedical researchers and scientists. They are not meant to cater to the needs of an individual patient. The web portal or any content(s) therein is neither designed to support, nor replace, the relationship that exists between a patient/site visitor and his/her physician. Your use of the WebmedCentral site and its contents is entirely at your own risk. We do not take any responsibility for any harm that you may suffer or inflict on a third person by following the contents of this website.